

# Phylogenetic Analysis, Cloning and Expression of Delta Amastin Encoding Gene from *Leishmania major* as a Vaccine Candidate

Zahra Yadegari<sup>1</sup>, Mohamad Reza Razavi<sup>2,3\*</sup>, Mitra Salehi<sup>1</sup>

<sup>1</sup>Department of biological science, North Tehran Branch, Islamic Azad University, Tehran, Iran. <sup>2</sup>Department of Parasitology, Pasteur Institute of Iran, Tehran, Iran. <sup>3</sup>Microbiology Research Center, Pasteur Institute of Iran, Tehran, Iran.

## ARTICLE INFO

### Research Article

VacRes, 2022

Vol. 9, No.1, 63- 72

Received: October 21, 2022

Accepted: November 17, 2022

Pasteur Institute of Iran

### \*Corresponding Author

Mohamad Reza Razavi;

Department of Parasitology, Pasteur

Institute of Iran, 69 Pasteur Ave., Tehran

1316943551, Iran.

Tel/Fax: +98 2164112257

Email: mrrazavi@yahoo.com

**KEYWORDS:** *Leishmania*, Amastin, Recombinant, Vaccine

## ABSTRACT

**Introduction:** Amastin is a surface glycoprotein in *Leishmania* species and one of the most important vaccine candidates due to its involvement in pathogenesis and being an essential virulence factor for parasite replication within the mammalian host cells. There are more than 60 copies of Amastin gene per genome of the parasite. **Methods:** Following phylogenetic analysis, a selected *Amastin* sequence was optimized and cloned with signal peptide in *Escherichia coli*. The recombinant protein was evaluated by SDS-PAGE and a specific antibody by Western blotting. **Results:** Among the *Amastin* sequences within different chromosomes of *Leishmania major*, the main type known as delta Amastin (primary distributed on chromosome 34) could be expressed in *E. coli* host and was confirmed by SDS-PAGE and Western blotting. **Conclusion:** Due to its copy number and evolutionary conservation and its role in pathogenesis,  $\delta$ -Amastin is considered as an important vaccine candidate against leishmaniasis which could be expressed as a recombinant protein in *E. coli*.

## INTRODUCTION

Leishmaniasis is a tropical disease induced by a vector borne parasite of *Leishmania* genus and is an endemic disease within 102 countries and five continents. Approximately, 350 million people are at risk of the infection while 15 million of them are known as infected cases reported worldwide with 1.5 to 2 million new cases and 70,000 per year [1]. Cutaneous Leishmaniasis (CL) is endemic in tropical and subtropical countries and approximately 90% of the cutaneous forms of the disease have been reported in the Middle East [2]. *Leishmania* parasite exhibits an intracellular amastigote form inside macrophages of the vertebrate host and is adapted to resist anti-parasitic activities such as acidic conditions, such as nitric oxide and reactive oxygen species and can also persist against the adapted immune responses [3].

Amastins are a vast family of transmembrane glycoproteins conserved among the *Trypanosomatids* which is mainly expressed during the amastigote form of the parasite [4-6]. Amastin of *Leishmania* amastigotes contains 11 amino acids at positions of 52-62 which are highly conserved and unique to *Trypanosomatid* protozoa [7]. It has been shown that Amastin proteins contain 4 hydrophobic transmembrane domains with two serine- and threonine-rich extracellular domains at N and C termini facing the cytosol. Comparative sequence analyses of *Amastin* genes have shown that the hydrophilic extracellular domain contains a conserved Amastin signature (i.e., C-[IVLYF]-[LF]-[WF]-G-X-[KRQ]-X-[DENT]-C) [8, 9].

Amastins are divided into 4 subfamilies of  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$  [5]. It is believed that  $\delta$ -Amastin subfamily is specific to *Leishmania* and none of these sequences were identified in other *Trypanosomatids*, neither by phylogenetic analysis nor by similarity-based protein clustering approach. These genes were developed from ancestral proto- $\delta$ -*Amastin* genes [10]. The Amastin family of proteins may be involved in transport of ions and protons across the membranes along with the regulation of cytoplasmic pH [11, 12]. Being a surface glycoprotein, Amastin can induce the immune system toward a Th1-like immune response [13]. Stober et al. have revealed that Amastin proteins are very immunogenic in mice at their signature sequence (i.e., CITLWGLKTDC) [14].

Approximately, 63 copies of  $\delta$ -Amastin have been found in the genome of *L. major*. The phylogenetic analysis of the Amastin sequence subfamily will reveal the main type of Amastin in the protozoa which involves in the adaptation of the parasite at the intracellular niche of the vertebrate host. It is envisaged that the selection and codon optimization of  $\delta$ -*Amastin* sequence may lead to efficient expression of this immunogenic protein in *E. coli*. This protein can be further investigated as a vaccine candidate for eliciting effective immune response and protection against Leishmaniasis.

## MATERIALS AND METHODS

### Phylogenetic Tree Reconstruction

The nucleotide sequences of Amastin gene were searched from GenBank, and were used based on LmajF.34.1960 *L. major* Friedlin strain putative Amastin-like surface protein partial mRNA 1-579 nucleotides, the aligned sequences were selected and used for phylogenetic tree construction. The phylogenetic relationships among the isolates were achieved from the phylogenetic tree reconstruction by the Neighbor Joining (NJ) method using MEGA 7.0 with assumption setting and the sure of the internal branches was tested with 1,000 bootstrap replications.

### Sequences Data of Amastin Gene of Kinetoplastida and *L. major* species.

The sequence of Amastin in *L. major* Friedlin strain LmajF.34.1960 (XM.001686241) was utilized for blast search and the sequences of Amastin gene sequences in order of Kinetoplastida including *Trypanosoma cruzi*, *Crithidia*, *Leishmania donovani*, *Leishmania infantum*, *Leishmania mexicana*, *Leishmania braziliensis* strains were used for tracing the phylogenetic tree construction. The phylogenetic association were analyzed by using the following sequences submitted to the GenBank, *Leishmania donovani*: MHOM/IN/1983/AG83 (ACCESSION CP019536), *Leishmania donovani* (ACCESSION CP022645.1), *Leishmania donovani*: LdBPK.080710 (ACCESSION FR799595.2), *Leishmania donovani*: LDHU3.24.1550 (ACCESSION LR812644.1), *Leishmania infantum*: LinJ.08.0680 (ACCESSION NC.009392.2), *Leishmania infantum*: LinJ.30.0930 (ACCESSION XM.001466901.1), *Leishmania infantum*: LinJ.24.1300 (ACCESSION FR796456.1), *Leishmania infantum*: LinJ.28.1510 (ACCESSION FR796460.1), *Leishmania mexicana*: LmxM.29.0850 (ACCESSION FR799582.1), *Leishmania mexicana*: LmxM.24.1270 (ACCESSION FR799577.1), *Leishmania mexicana*: LmxM.28.1400 (ACCESSION FR799581.1), *Leishmania mexicana*: LmxM MHOM/GT/2001/U1103 (ACCESSION XM.003886514.1), *Trypanosoma cruzi*: Tc00.1047053509051.20 (ACCESSION XM\_801067.1), *Trypanosoma cruzi*: Tc00.1047053507485.150 (ACCESSION XM\_807298.1), *Leishmania braziliensis*: LBRM.08.0680 (ACCESSION NC\_009300.2), *Leishmania braziliensis*: LbrM.24.1280 (ACCESSION NC\_009316.2), *Leishmania braziliensis*: LbrM.28.1550 (ACCESSION XM\_003723148.1), *Leishmania braziliensis*: LbrM.30.0980 (ACCESSION XM\_001566681.1), a-amastin [*Crithidia* sp. ATCC 30255]: CDFL9H15.10 (ACCESSION ACS87875.1), b-amastin [*Crithidia* sp. ATCC 30255]: CDFL7M16.03 (ACCESSION ACS87856.1), d-amastin, partial [*Crithidia* sp. ATCC 30255]: CDFL2E16.01 (ACCESSION ACS87802.1), g-amastin [*Crithidia* sp. ATCC 30255]: CDFL6B24.01 (ACCESSION GQ153665.1)

The sequence of Amastin in *L. major* Friedlin strain LmajF.34.1960 (XM.001686241) was used to find the homologous sequences in genome of *L. major* Friedlin strain. The phylogenetic relationship were analyzed by using the following sequences ( LmajF.08.0670 ) ACCESSION XM.001681044.1 ( , LmajF.08.0680 ) ACCESSION XM.001681045.1 ( , LmajF.08.0690 ) ACCESSION XM.001681046.1 ( , LmajF.08.0700 ) ACCESSION XM.001681047.1 ( , LmajF.08.0710 ) ACCESSION

XM.001681048.1 ( , LmajF.08.0720 ) ACCESSION  
 XM.001681049.1 ( , LmajF.08.0730 ) ACCESSION  
 XM.001681050.1 ( , LmajF.08.0740 ) ACCESSION  
 XM.001681051.1 ( , LmajF.08.0750 ) ACCESSION  
 XM.001681052.1 ( , LmajF.08\_0760 ) ACCESSION  
 XP.001681105 ( , LmajF.08.0770 ) ACCESSION  
 XM.001681054.1 ( , LmajF.08.0800 ) ACCESSION  
 XM.001681057.1 ( , LmajF.08.0810 ) ACCESSION  
 XM.001681058.1 ( , LmajF.08.0820 ) ACCESSION  
 XM.001681059.1 ( , LmajF.08.0830 ) ACCESSION  
 XM.001681060.1 ( , LmajF.08.0840 ) ACCESSION  
 XM.001681061.1 ( , LmajF.08.0850 ) ACCESSION  
 XM.001681062.1 ( , LmajF.08.0640 ) ACCESSION  
 XM.001681041.1 ( , LmajF.10.0220 ) ACCESSION  
 XM.001681302.1 ( , LmajF.10.0230 ) ACCESSION  
 XM.001681303.1 ( , LmajF.16.0490 ) ACCESSION  
 XM.001682075.1 ( , LmajF.24.1250 ) ACCESSION  
 XP.001683641 ( , LmajF.24.1260 ) ACCESSION  
 XM.001683590.1 ( , LmajF.24.1270 ) ACCESSION  
 XP.001683643 ( , LmajF.24.1280 ) ACCESSION  
 XM.001683592.1 ( , LmajF.27.0540 ) ACCESSION  
 XM.003721812.1 ( , LmajF.28.1120 ) ACCESSION  
 XM.001684342.1 ( , LmajF.28.1130 ) ACCESSION  
 XM.001684343.1 ( , LmajF.28.1390 ) ACCESSION  
 XM.001684369.1 ( , LmajF.28.1400 ) ACCESSION  
 XP.001684422 ( , LmajF.30.0850 ) ACCESSION  
 XP.001684680.1 ( , LmajF.30.0860 ) ACCESSION  
 XP.001684681 ( , LmajF.30.0870 ) ACCESSION  
 XP.001684682 ( , LmajF.31.0450 ) ACCESSION  
 XP.001685030 ( , LmajF.34.0500 ) ACCESSION  
 XP.001686141 ( , LmajF.34.0960 ) ACCESSION  
 XM.001686137 ( , LmajF.34.0970 ) ACCESSION  
 XM.001686138.1 ( , LmajF.34.0980 ) ACCESSION  
 XM.001686139.1 ( , LmajF.34.1080 ) ACCESSION  
 XM.001686149.1 ( , LmajF.34.1560 ) ACCESSION  
 XM.001686201.1 ( , LmajF.34.1580 ) ACCESSION  
 XM.001686203.1 ( , LmajF.34.1600 ) ACCESSION  
 XM.001686205.1 ( , LmajF.34.1620 ) ACCESSION  
 XM.001686207.1 ( , LmajF.34.1640 ) ACCESSION  
 XM.001686209.1 ( , LmajF.34.1660 ) ACCESSION  
 XM.001686211.1 ( , LmajF.34.1680 ) ACCESSION  
 XM.001686213.1 ( , LmajF.34\_1700 ) ACCESSION  
 XM.001686215.1 ( , LmajF.34.1720 ) ACCESSION  
 XM.001686217.1 ( , LmajF.34.1740 ) ACCESSION  
 XM.001686219.1 ( , LmajF.34.1760 ) ACCESSION  
 XM.001686219.1 ( , LmajF.34.1780 ) ACCESSION  
 XM.001686223.1 ( , LmajF.34.1800 ) ACCESSION  
 XM.001686225.1 ( , LmajF.34.1820 ) ACCESSION  
 XM.001686227.1 ( , LmajF.34.1840 ) ACCESSION  
 XM.001686229 ( , LmajF.34.1860 ) ACCESSION  
 XM.001686231.1 ( , LmajF.34.1880 ) ACCESSION  
 XM.001686233.1 ( , LmajF.34.1900 ) ACCESSION  
 XM.001686235.1 ( , LmajF.34.1920 ) ACCESSION  
 XM.001686237.1 ( , LmajF.34.1940 ) ACCESSION  
 XM.001686239.1 ( , LmajF.34.1960 ) ACCESSION  
 XM.001686241 ( , LmajF.34.1980 ) ACCESSION  
 XM.001686243.1 ( , LmajF.36.1270 ) ACCESSION  
 XM.001686647.1 ( , LmajF.36.4140 ) ACCESSION  
 XM.001686945.1.

### Analysis of Amino Acid Sequences of Amastin

The amino acid sequences of Amastin, Amastin-like and putative Amastin proteins of *L. major* Friedlin strain were

aligned using MEGA7 software (Molecular Evolutionary Genetics Analysis, version 7.0 for bigger datasets) [15]. The conserved amino acid sequence parts similar to Amastin of LmjF.34.1960 with 192 amino acids were determined by default setting.

### Bacterial strains, Plasmids, and Chemicals

*E. coli* DH5 $\alpha$  was used as the cloning host and *E. coli* BL21 (DE3) strain as the expression host. For cultivation of the cloning and the expression hosts, ampicillin (100  $\mu$ g/ml) and kanamycin (33  $\mu$ g/ml) antibiotics were added to the culture medium, respectively. Plasmid pUC19 was used as the cloning and pET28 was utilized as the expression vectors. When required, for transformation of the cloning vector in competent cells, *E. coli* DH5 $\alpha$  strain competent was prepared by a 2:100 overnight culture transfer in fresh LB broth (37 °C, at 180 rpm, 3 h). The culture medium was transferred on ice for 1 h. After centrifuging 1.5 ml of the culture medium (5 min, 5000 rpm, 10 min, 4 °C), the pellet was resuspended in 0.5 ml of 100 mM CaCl<sub>2</sub> and put on ice for 45 min. The cells were washed again and were resuspended in 100  $\mu$ l of 100mM CaCl<sub>2</sub>. For transformation of cells, approximately 0.5  $\mu$ g of plasmid was mixed with 50  $\mu$ l suspension of competent cells and incubated on ice for 45 min. The cells were transferred on hot plate at 42 °C for 90 s, and then were put on ice. After adding 500  $\mu$ l LB broth to the pellet and incubation at 37 °C for 45 min, 150  $\mu$ l samples with and without the construct were cultured on LB agar plate containing the antibiotic and incubated 37 °C for 16-20 h. The transformed clones were selected and cultured in 4 ml LB medium containing the antibiotic and incubated for 16-24 h at 37 °C. The plasmid was purified with Favorgen kit (Yektatajiz Azma, Iran) according to the manufacturer's instruction and analyzed by restriction map preparation.

### Construction of the Expression Vector with a Signal Peptide of *E. coli*

The pUC57 cloning vector containing nucleotide optimized sequence (Pishgam Biotech Company, Iran) had a  $\delta$ -Amastin sequence containing the *Hind* III restriction sequences at both ends. After transformation and plasmid purification, the vector was digested with *Hind* III and the digested sequence was extracted from agarose gel by GeneAll kit (GeneAll Biotechnology, South Korea).

The type-1 fimbrial subunit of *E. coli* is encoded by *fimA* gene. To design primers for amplification of *fimA* gene signal peptide, type 1 fimbriae major subunit of *E. coli* strain K-12, substrain MG1655 was selected (NCBI GeneID: 948838; 549 bp). Using Gene Runner software, the primers were designed for the signal peptide part of this gene, and *Nde*I restriction site was placed in the forward primer (GTTTTTTGAAAGGAAAGCCATATG), and *Hind* III restriction site was placed in the reverse primer (AAAAGCTTGGTCCACCATTAAC). The restriction sites are shown in bold in the sequence. The amplification reaction for the signal peptide was conducted using the designed primers and the genomic DNA of *E. coli* which was purified and used as the template. The reaction began with an initial denaturation step at 96 °C for 3 min and 30 cycles of 45 s at 96 °C, 45 s at 56 °C, and 60 s at 72 °C with a final extension at 72 °C for 5 min. After digestion and purification of the amplicon, it was ligated with a pET28 plasmid digested with the same restriction enzymes. In order to transform of *E. coli* DH5 $\alpha$  competent cells, 5  $\mu$ l of the ligation mix were used. The transformed clones were confirmed by plasmid purification and restriction

digestion. The clones containing *fimA* sequence were used for the next step.

Both the purified pET28 plasmid (with the signal peptide sequence) from the previous step and pUC57 vector containing  $\delta$ -Amastin sequence were digested with *Hind* III enzyme. Both ends of the amplified DNA fragment cleavage site and the ends of the vector used in this study were sticky. A molar ratio of 1:2 vector: insert gel purified DNA was used to prepare the ligation mixture (4 °C, 16 h) with T4 DNA Ligase (Thermo Fisher Scientific Inc.). The product was added to *E. coli* DH5 $\alpha$  competent cells. The transformed colonies were identified by purified plasmid size comparison and enzyme-digestion using *Hind* III and *Nde* I enzymes. The selected transformants of the cloning host were used for plasmid purification and the purified product was transformed to the expression host BL21 DE3. The resulting clones were examined for the expression of  $\delta$ -Amastin.

### Recombinant $\delta$ -Amastin Protein Expression

A clone of the transformed plasmid in the BL21 DE3 expression host was cultured in LB medium containing Kanamycin 33  $\mu$ g/ml and incubated for 16-24 hours at 37 °C. Subculture was then performed in the fresh medium and shacked for 4 hours at 37 °C at 180 rpm. After reaching the optical absorption of the culture medium to 0.6 at a wavelength of 600 nm, equal volume of fresh medium with kanamycin 33  $\mu$ g/ml was added to culture medium. To the final volume of the culture, IPTG was tested at different final concentrations (0.2, 0.4, 0.6 and 0.8 mM). The medium was incubated at 25 °C or 37 °C and shacked at 180 rpm for 20 h. The cells were harvested by centrifugation at 5000 rpm for 5 minutes.

### Confirmation of $\delta$ -Amastin Protein Expression by SDS-PAGE, Dot Blot and Western Blotting

A 500  $\mu$ l culture medium of cells harboring the construct were harvested and lysed in 40  $\mu$ l of 50 mM Tris-HCl (pH 6.8) by vortexing. After adding the sample buffer, the mixture was boiled and was incubated on ice for 10 min. The sample was centrifuged at 6000 rpm (4°C, 10 min) and loaded on 12% polyacrylamide gel. The gel was stained with 0.05% Coomassie brilliant blue for 40 min and then destained (5% acetic acid, 25% methanol v/v).

To investigate the expression of the target protein in the cell lysate, the dot blot method was performed. Fifty  $\mu$ l of sample pellet was dissolved in 50 mM Tris solution (pH 6.8) and vortexed. The cell lysate was centrifuged, poured on the nitrocellulose membrane and dried. The nitrocellulose membrane was incubated in the blocking solution (2% skimmed milk powder without fat in TBST, NaCl 250 mM, Tris Base 10 mM, and pH 7.5) for 1 h at room temperature on the shaker. The membrane was washed 3 times with Washing Buffer (TBST; every time for 10 min on shaker at room temperature). The membrane was placed in the primary antibody solution (anti-His Tag Human-Mouse hybrid monoclonal diluted 1:750 in TBST; Sina Biotch Company, Iran), for 1 h on shaker at room temperature and washed again. In the next step, the membrane was soaked in a secondary antibody solution (Sheep anti-human IgG HRP conjugated polyclonal antibody diluted 1:1000 in TBST; Sina Biotch Company, Iran), for 1 h at room temperature on the shaker. The antibody was washed off 3 times with TBST. Finally, 10 mg of DAB, 20 ml of TBST and 1  $\mu$ l of H<sub>2</sub>O<sub>2</sub> 30% were mixed and poured on the nitrocellulose membrane and kept in the dark for 15 to 30 min. After the dot appeared, the membrane was washed with distilled water.

For Western blotting, the lysate of the protein sample was prepared as above. The protein concentration was determined with spectrophotometry based on absorbance at 280 nm. The protein samples (1 to 2 mg) were gel-loaded and run by SDS-PAGE. The gel-separated proteins bands were transferred to a nitrocellulose membrane by electro-transfer buffer (25 mM Tris, 192 mM glycine, pH 8.3, 20% v/v methanol; 1 h, 80 V). The membrane was washed with distilled water. The nitrocellulose membrane was incubated in the blocking buffer for 1 h and followed as described above for the dot blot procedure.


## RESULTS

### Phylogenetic Relationship of Amastin Gene within Kinetoplastida and *L. major* Species

Amastins have homology in species and different strain of *Leishmania*. It has four subfamilies  $\alpha$ ,  $\beta$ ,  $\delta$  and  $\gamma$ . In this study, the similarity of LmjF.34.1960 sequence was searched with the selected *Amastin* sequences within the order of Kinetoplastida. According to the constructed tree as shown in Fig. 1, the sequences of *L. donovani*: LDHU3.24.1550, *L. infantum*: LinJ.24.1300, g-amastin [*Crithidia* sp. ATCC 30255]: CDFL6B24.01, *L. mexicana*: LmxM.24.1270, *L. major*:

LmjF.24.1270, *L. braziliensis*: LbrM.24.1280 and *L. major* and LmjF.24.1280 include in the subfamily  $\gamma$ . The subfamily  $\alpha$  contains the Amastin sequences a-amastin [*Crithidia* sp. ATCC 30255]: CDFL9H15.10, *L. braziliensis*: LbrM.28.1550, *L. mexicana*: LmxM.28.1400, *L. major*: LmjF.28.1400, *L. infantum*: LinJ.28.1510 and *L. donovani*: MHOM/IN/1983/AG83.

In the clade of subfamily  $\delta$ , *L. infantum*: LinJ.08.0680, *L. donovani*: LdBPK.08.0710, *L. mexicana*: LmxM.29.0850, *L. mexicana*: LmxM.33.1900.1, d-amastin, partial [*Crithidia* sp. ATCC 30255]: CDFL2E16.01, *T. cruzi*: Tc00.1047053507485.150, *L. major*: LmjF.08.0760, *L. major*: LmjF.08.0810, *L. major*: LmjF.31.0450, *L. major*: LmjF.34.1840, *L. major*: LmjF.34.1860 and LmjF.34.1960 in *L. major* Friedlin strain are seen. Finally, the sequences of *L. major*: LmjF.30.0860, *L. braziliensis*: LbrM.08.0680, *L. braziliensis*: LbrM.30.0980, *L. infantum*: LinJ.30.0930, *L. major*: LmjF.30.0870, b-amastin [*Crithidia* sp. ATCC 30255]: CDFL7M16\_03, *T. cruzi*: Tc00.1047053509051.20, *L. donovani*: CP022645.1 are within the subfamily  $\beta$ . The result revealed that the LmjF.34.1960 in *L. major* Friedlin strain is in the subfamily  $\delta$ . There are not any *Amastin* sequences of *T. cruzi* related to the  $\alpha$  and  $\gamma$  subfamilies.



**Fig. 1.** Phylogenetic relationship of sequences  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ -Amastin subfamilies of Kinetoplastida (*T. cruzi*, *Crithidia* and *Leishmania* species) based on LmjF.34.1960 *L. major* Friedlin strain putative Amastin-like surface protein partial mRNA 1-579 nucleotides. *Amastin* sequences of different species are divided in four clades. The evolutionary relationship was prepared using the Neighbor-Joining method, using MEGA 7.0 by default setting and the reliability of the internal branches was tested by 1,000 bootstrap replications.

### Analysis of Amastin Amino Acid Sequences in *L. major* Friedlin strain and their Phylogenic Relationship

There are 63 sequences of Amastin genes located in 9 different chromosomes in *L. major* Friedlin strain. They are classified in four amastin subfamilies,  $\alpha$ ,  $\beta$ ,  $\delta$  and  $\gamma$ - amastin. There are 44 numbers  $\delta$ -Amastin on chromosome 8, 31, 34, 36; 2 proto- $\delta$  ( $p\delta$ ) Amastin on chromosome 34; 2  $\alpha$ -Amastin genes on chromosome 28; 4  $\gamma$  amastin on chromosome 24 and 11  $\beta$  amastin on chromosome 8,10,16, 27, 28, 30, 36 in *L. major* Friedlin strain. The prominent type of amastin sequence in the genome of *L. major* Friedlin strain is  $\delta$  -amastin which including more than 73 percent of amastin gene copies followed by  $\beta$ -amastin copies with more than 17 percent. The  $\alpha$  and  $\gamma$ -


amastin types form about 10 percent of gene copy numbers. (Table 1).

In *L. major* Friedlin strain, the conserved amino acids sequences of Amastins are 11 amino acids (C-[IVLYF]-[TS]-[LF]-[WF]-G-X-[KRQ]-X-[DENT]-C), 2 amino acid G [FY] in position 40-50 and 152-153 according to the LmjF.34.1960 sequence, respectively. The conserved 11 amino acids segment is absent in 8 sequences of 63 amastin protein copies in *L. major* as follow LmjF.10.0220, LmjF.10.0230, LmjF.16.0490, LmjF.27.0540, LmjF.08.0640, LmjF.28.1120, LmjF.28.1130 LmjF.36.4140. The conserved motif of G [FY] at position 152-153 is absent in three copies, LmjF.16.0490, LmjF.27.0540, LmjF.36.4140.


**Table 1.** Comparison of 63 sequences of amastin gene copies located on 9 different chromosomes in *L. major* Friedlin strain with their accession number, amastin type, conserved sequence domain and related chromosome. the conserved amino acids sequences are 11 amino acids (C-[IVLYF]-[TS]-[LF]-[WF]-G-X-[KRQ]-X-[DENT]-C), 2 amino acid G[FY] in position 40-50 and 152-153 of LmjF.34.1960, respectively.

Gene	Accession number	Conserve sequences of protein		Chromosome	Type
<a href="#">LmajF.08.0670</a>	XM.001681044.1	CITLWGAKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0680</a>	XM.001681045.1	CITLWGAKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0690</a>	XM.001681046.1	CITLWGAKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0700</a>	XM.001681047.1	CITLWGLKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0710</a>	XM.001681048.1	CITLWGAKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0720</a>	XM.001681049.1	CITLWGLKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0730</a>	XM.001681050.1	CITLWGAKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0740</a>	XM.001681051.1	CITLWGLKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0750</a>	XM.001681052.1	CITLWGAKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0760</a>	XP.001681105	CITLWGLKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0770</a>	XM.001681054.1	CITLWGLKTDC	GF	8	$\delta$
<a href="#">LmajF.08.0800</a>	XM.001681057.1	CLTLWGGKEEC	GF	8	$\delta$
<a href="#">LmajF.08.0810</a>	XM.001681058.1	CLTLWGYKSEC	GF	8	$\delta$
<a href="#">LmajF.08.0820</a>	XM.001681059.1	CLTLWGGKEEC	GF	8	$\delta$
<a href="#">LmajF.08.0830</a>	XM.001681060.1	CLTLWGGKEEC	GF	8	$\delta$
<a href="#">LmajF.08.0840</a>	XM.001681061.1	CLTLWGGKEEC	GF	8	$\delta$
<a href="#">LmajF.08.0850</a>	XM.001681062.1	CLTLWGYKSEC	GF	8	$\delta$
<a href="#">LmajF.08.0640</a>	XM.001681041.1	_____	_____	8	$\beta$
<a href="#">LmajF.10.0220</a>	XM.001681302.1	_____	_____	10	$\beta$
<a href="#">LmajF.10.0230</a>	XM.001681303.1	_____	_____	10	$\beta$
<a href="#">LmajF.16.0490</a>	XM.001682075.1	_____	GF	16	$\beta$
<a href="#">LmajF.24.1250</a>	XP.001683641	CYTMWGYRKF	GF	24	$\gamma$
<a href="#">LmajF.24.1260</a>	XM.001683590.1	CYTFIGYKTEC	GF	24	$\gamma$
<a href="#">LmajF.24.1270</a>	XP.001683643	CYTFWGMKSDC	GF	24	$\gamma$
<a href="#">LmajF.24.1280</a>	XM.001683592.1	CYTMWGYRKFC	GF	24	$\gamma$
<a href="#">LmajF.27.0540</a>	XM.003721812.1	_____	GF	27	$\beta$
<a href="#">LmajF.28.1120</a>	XM.001684342.1	_____	GF	28	$\beta$
<a href="#">LmajF.28.1130</a>	XM.001684343.1	_____	GF	28	$\beta$
<a href="#">LmajF.28.1390</a>	XM.001684369.1	CYTFWGYRANC	GF	28	$\alpha$


<a href="#">LmajF.28.1400</a>	XP.001684422	CFTYWGKYKDNC	GF	28	$\alpha$
<a href="#">LmajF.30.0850</a>	XP.001684680.1	CVTVWGLKNDC	GY	30	$\beta$
<a href="#">LmajF.30.0860</a>	XP.001684681	CVTVWGLKNDC	GY	30	$\beta$
<a href="#">LmajF.30.0870</a>	XP.001684682	CFTLWGLHSDC	GY	30	$\beta$
<a href="#">LmajF.31.0450</a>	XP.001685030	CLTLWGEKLDC	GF	31	$\delta$
<a href="#">LmajF.34.0500</a>	XP.001686141	CITLWGWKSR	GF	34	$\delta$
<a href="#">LmajF.34.0960</a>	XM.001686137	CVTLWGAKLGC	GF	34	$\delta$
<a href="#">LmajF.34.0970</a>	XM.001686138.1	CLTIWGFKDKC	GF	34	$p\delta$
<a href="#">LmajF.34.0980</a>	XM.001686139.1	CVSLWGIRDRC	GF	34	$p\delta$
<a href="#">LmajF.34.1080</a>	XM.001686149.1	CVTLWGAKLGC	GF	34	$\delta$
<a href="#">LmajF.34.1560</a>	XM.001686201.1	CITLWGEFTC	GF	34	$\delta$
<a href="#">LmajF.34.1580</a>	XM.001686203.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1600</a>	XM.001686205.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1620</a>	XM.001686207.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1640</a>	XM.001686209.1	CITLWGLRLTC	GF	34	$\delta$
<a href="#">LmajF.34.1660</a>	XM.001686211.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1680</a>	XM.001686213.1	CLTLFGLKFDC	GF	34	$\delta$
<a href="#">LmajF.34.1700</a>	XM.001686215.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1720</a>	XM.001686217.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1740</a>	XM.001686219.1	CITLWGERFTC	GF	34	$\delta$
<a href="#">LmajF.34.1760</a>	XM.001686219.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1780</a>	XM.001686223.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1800</a>	XM.001686225.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1820</a>	XM.001686227.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1840</a>	XM.001686229	CITLFGVKVDC	GF	34	$\delta$
<a href="#">LmajF.34.1860</a>	XM.001686231.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1880</a>	XM.001686233.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1900</a>	XM.001686235.1	CITLWGERFTC	GF	34	$\delta$
<a href="#">LmajF.34.1920</a>	XM.001686237.1	CITLWGDKLAC	GF	34	$\delta$
<a href="#">LmajF.34.1940</a>	XM.001686239.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1960</a>	XM.001686241	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.34.1980</a>	XM.001686243.1	CLTLFGFKLDC	GF	34	$\delta$
<a href="#">LmajF.36.1270</a>	XM.001686647.1	CITLWGLRLTC	GF	36	$\delta$
<a href="#">LmajF.36.4140</a>	XM.001686945.1	—————	GY	36	$\delta$



**Fig. 2.** Phylogenetic relationship of sequences  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $\delta$ -Amastin subfamilies of of *L. major* Friedlin strain based on LmjF.34.1960 *L. major* Friedlin strain putative Amastin-like surface protein partial mRNA 1-579 nucleotides. Amastin sequences of different copies are divided mainly in the  $\delta$ -Amastin clade, the minority, less than 30% of the copies form the other 3 clads, namely  $\alpha$ ,  $\beta$  and  $\gamma$ -Amastin. The evolutionary relationship was prepared using the Neighbor-Joining method, using MEGA 7.0 by default setting and the reliability of the internal branches was tested by 1,000 bootstrap replications.



**Fig. 3.** The PCR product amplified with two specific primers for the *fimA* gene signal peptide in 1% agarose gel and 100 bp DNA Ladder (Vazyme). Lane M: DNA Ladder, lane 1 to 3: the PCR product amplification of the signal peptide of the *fimA* gene in the genomic DNA of *E. coli*.




**Fig. 4.** The pET28 expression construct containing *fla* gene signal peptide and amastin coding sequence digestion with *Hind* III enzyme and without enzymatic cleavage. Lane M: 100 bp DNA Ladder (Vazyme), lane 1: The construct without enzymatic cleavage, lane 2: The construct with *Hind* III enzyme digestion.

#### Recombinant $\delta$ -Amastin Expression


The expression of recombinant  $\delta$ -Amastin protein with an expected molecular weight of 37 kDa was examined at 0.2, 0.4, 0.6, and 0.8 mM IPTG at 25°C and 37 °C temperatures. The results indicated that the highest amastin expression was achieved with an IPTG concentration of 0.8 mM and at 25°C (Fig. 5A). The expression at 25°C was also assayed using pellets prepared from the culture medium with 0.8 mM IPTG protein expression at 12, 24, and 36 h after the induction (Fig. 5B).

#### Confirmation of $\delta$ -Amastin Expression by Dot Blot and Western Blotting

The recombinant expressed protein was confirmed using the host cell lysate (0.8 mM IPTG induction at 25 °C for 24 h) by a dot blot procedure. The reaction of recombinant  $\delta$ -Amastin with a specific anti-His tag antibody was revealed by an intense colored dot compared to the negative control on the membrane (Fig. 6 A). For confirmation of the expressed protein molecular weight, the cell lysate was analyzed by Western blotting which detected an approximately 37 kDa band using a specific anti-His tag antibody (Fig. 6 B).



**Fig. 5. (A)** Expression of recombinant  $\delta$ -Amastin with different IPTG concentrations (0.2 mM to 0.8 mM) at 25 °C . Lane M: Pre-stained Protein Molecular weight Marker (Sinaclon), Lane 1: uninduced negative control, Lane 2: induced at 25 °C with 0.2 mM IPTG, Lane 3: induced at 25 °C with 0.4 mM IPTG, Lane 4: induced at 25 °C with 0.6 mM IPTG, Lane 5: induced at 25 °C with 0.8 mM IPTG. **(B)** Investigation of Amastin expression at different time intervals after 0.8 mM IPTG induction at 25 °C. Lane M: Pre-stained Protein Molecular weight Marker (Biolegend, USA), Lane 1: uninduced negative control, Lane 2: 12 h after the induction, Lane 3: 24 h after the induction, Lane 4: 36 h after the induction of protein expression.



**Fig. 6.** Final Confirmation of recombinant  $\delta$ -Amastin expression by dot blot and Western blotting methods. **A)** Dot blot: Lane 1: negative control, Lane 2: Recombinant  $\delta$ -Amastin test sample. **B)** Western blotting procedure, Lane M: Pre-stained Protein Molecular Weight Marker (Sinaclon, Iran), Lane 1: Recombinant  $\delta$ -Amastin, Lane 2: Negative control.

## DISCUSSION

Leishmaniasis is a neglected vector-borne disease in tropical and subtropical areas with increased prevalence during the last decade. The infection is correlated to the socio-economic conditions in the endemic foci. At the present, the control measurements against leishmaniasis are mainly relied on controlling the vectors and the reservoirs of the parasite. As a result, the sensitive population who remain at the infected areas can increase the risk of infection in the subsequent years in those areas. Hence, studying the antigenic molecules as vaccine candidates which are capable of eliciting effective immune responses in human population and protection against leishmaniasis in the infected areas is a public health priority.

Amastin surface proteins are encoded by a broad family of genes known so far in 45 species of *Leishmania*. All of these genes have a common structure and have an extracellular domain with 11 highly conserved amino acids. Most of these genes are expressed in the amastigote stage of the parasite and are present upstream of 3'-UTR. A fragment of 450 nucleotides is conserved in the region that regulates the gene expression at the transcriptional level, and possibly other mechanisms may be involved in controlling or regulating their transcription in macrophages [4]. Another study has shown that a 450-nucleotide region responds to heat shock by initiating the transcription of Amastin mRNA in *Leishmania*. Upon entering the host cell, it has also been shown that another 100-nucleotide region in the 3'-end of Amastin mRNAs increase the translation of this protein in response to temperature. In addition, the acidic pH present in phagolysosomes and macrophages has been shown to produce large amounts of Amastin mRNA, which is independent of the 450-nucleotide and the 100-nucleotide fragments. Post-transcriptional regulation of Amastin mRNA in *Leishmania* is complex and various mechanisms are involved in its stability and translation [16]. The expression of this protein is induced at an acidic pH. Moreover, the fact that Amastin encoding gene is conserved in all species of *Leishmania*, is of particular importance. Due to its immunogenicity, Amastin has been used for both diagnosis and immunization [7]. For instance, in a study by Vale *et al.*, Amastin of *L. infantum* was evaluated for disease diagnosis in dogs and humans. The results indicated high sensitivity of these tests for diagnosis of the disease when used simultaneously in dogs and humans [11].

In order to obtain an appropriate antigen for immunization against *L. major* infection in this study, the conserved and common form of *Amastin* gene was cloned into a pET expression vector and was expressed in *E. coli* and the results showed a proper expression of the antigen. By examining the suitable temperature conditions for the optimal expression of the recombinant protein, 25°C was found to be the best temperature for the expression. The likely causes of this are the slow metabolism and decreased secretion of toxic metabolites in the culture medium at this temperature. The IPTG concentrations in range of 0.5 and 1 mM IPTG indicated that 0.8 M concentration was the most appropriate IPTG concentration for the optimal expression of Amastin in *E. coli*. The expression of the recombinant protein at different times after the induction indicated that the expression began at approximately 9 hours after the induction and increased to up to approximately 24 hours with no increase in the amount of protein after this time point.

Previously, Zhang and colleagues have revealed multiepitope vaccine composed of KMP-11, Amastin and GP63 proteins is protective in BALB/c mice. Following two

injections, the animals were challenged with *Leishmania* promastigotes. In the animal sera the antibodies specific to the epitopes and in spleen cells secreted cytokines CD3+, CD4+, CD8+ were detected by the ELISA technique. Moreover, the parasitic load was reduced in the GP63, KMP-11, and amastin treated groups to 89, 86, and 79 percent compared to the control group [22].

The amastin of *L. infantum* has been evaluated for immunogenicity and protection in BALB/c mice. It has been reported that at forty-five days after infection with *L. infantum* promastigotes, based on protein and parasite-specific IFN- $\gamma$ , IL-12, GM-CSF, and the IgG2a isotype antibody, a Th1 immune response occurs in vaccinated mice [23].

The phylogenetic relationship of Amastin protein in *L. major* indicated that the majority of Amastin gene copies are in the main form of  $\delta$ -Amastin which comprise more than 73 % of the copies. This form of Amastin is evolutionary one of the most conserved forms; therefore, has an important role in biology and virulence of the parasite, especially in the vertebrate host cells. The evaluation of the immunogenicity and efficacy of Amastin in animal models have revealed that it can elicits protective immune responses in animals against leishmaniasis; although, the type of the Amastins that were used in these studies are not well-specified [17]. The placement of an *E. coli* signal peptide upstream of the Amastin encoding gene transports the expressed protein into the bacterial cell membrane; thereby, it can reduce the concentration of Amastin within the cytoplasm during the induction of the expression by IPTG. This in turn, reduces the transcription and translation inhibition of the Amastin gene, resulting in continuous expression of the protein in the cell, in its natural form.

In this study, the total cellular protein amount was 50  $\mu\text{g/ml}$  of the culture medium while the expression of Amastin was evaluated as 0.5  $\mu\text{g/ml}$  of the culture medium. This indicated that the expression of the protein in the cell can increase up to approximately 1-2% of the total amount of the cellular protein at the best environmental conditions. Since in other studies, 5 to 50  $\mu\text{g}$  of this protein has been used for immunization [13, 18], the obtained concentration of the recombinant Amastin by the described method seems to be sufficient for immunization in future animal studies. Therefore, the next step is to prepare the appropriate dose and formulation of this recombinant protein to assess the outcome of using it for the immunization assays.

In conclusion, Amastin is one of the most important vaccine candidates for protection of sensitive population in the endemic areas against leishmaniasis. The phylogenetic analysis of Amastin surface glycoprotein protein family encoding sequences in *L. major* revealed that the abundant type is  $\delta$ -amastin which is widely distributed on chromosomes 34 and 8 of the parasite. The copy number and the evolutionary conservation of this prominent type emphasize its role in the pathogenesis and survival of the parasite in the vertebrate hosts. Due to the conserved nature of this virulence factor within the genus of *Leishmania* and *Trypanosomatidae* family, the cloning and expression of  $\delta$ -Amastin encoding sequence described in this study appears to be a promising mean to prepare an effective antigen for induction of protective immune response in future animal studies.

## ACKNOWLEDGMENT

The authors would like to thank Departments of Mycobacteriology as well as Hepatitis and HIV of Pasteur Institute of Iran for providing certain laboratory facilities. The authors would like to appreciate Ms. Mariam Amin for her supervision and laboratory assistance.

## CONFLICT OF INTEREST

The authors declare they have no conflict of interests.

## REFERENCES

- Alvar J, Vélez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. World Health Organization, WHO Leishmaniasis Control Team Leishmaniasis worldwide and global estimates of its incidence PLoS One. 2012;7(5):e35671.
- Razavi MR, Shirzadi MR, Mohebbi M, Yaghoobi-Ershadi MR, Vatandoost H, Shirzadi M et al. Human Cutaneous Leishmaniasis in Iran, Up to Date-2019. Journal of Arthropod-Borne Diseases. 2021;15(2):143.
- Chang K-P, Fish WR. Leishmania. In vitro cultivation of protozoan parasites. 2019:111-54.
- Butenko A, Hammond M, Field MC, Ginger ML, Yurchenko V, Lukeš J. Reductionist pathways for parasitism in euglenozoans? Expanded datasets provide new insights. Trends in parasitology. 2021;37(2):100-16.
- Jackson AP. The evolution of amastin surface glycoproteins in trypanosomatid parasites. Molecular biology and evolution. 2010;27(1):33-45.
- Liu Q, Lei J, Darby AC, Kadowaki T. Trypanosomatid parasite dynamically changes the transcriptome during infection and modifies honey bee physiology. Communications biology. 2020;3(1):1-8.
- de Paiva RMC, Grazielle-Silva V, Cardoso MS, Nakagaki BN, Mendonca-Neto RP, Canavaci AMC et al. Amastin knockdown in *Leishmania braziliensis* affects parasite-macrophage interaction and results in impaired viability of intracellular amastigotes. PLoS pathogens. 2015;11(12):e1005296.
- Rochette A, McNicoll F, Girard J, Breton M, Leblanc É, Bergeron MG et al. Characterization and developmental gene regulation of a large gene family encoding amastin surface proteins in *Leishmania* spp. Molecular and biochemical parasitology. 2005;140(2):205-20.
- Cerqueira GC, Bartholomeu DC, DaRocha WD, Hou L, Freitas-Silva DM, Machado CR et al. Sequence diversity and evolution of multigene families in *Trypanosoma cruzi*. Molecular and biochemical parasitology. 2008;157(1):65-72.
- Butenko A, Opperdoes FR, Flegontova O, Horák A, Hampl V, Keeling P et al. Evolution of metabolic capabilities and molecular features of diplomids, kinetoplastids, and euglenids. BMC biology. 2020;18(1):1-28.
- Vale DL, Dias DS, Machado AS, Ribeiro PA, Lage DP, Costa LE et al. Diagnostic evaluation of the amastin protein from *Leishmania infantum* in canine and human visceral leishmaniasis and immunogenicity in human cells derived from patients and healthy controls. Diagnostic Microbiology and Infectious Disease. 2019;95(2):134-43.
- Biswas B, Laha B, Sengupta P, Das K, Ghosh M. Characterization of a novel amastin-like surface protein (ALSP) of *Leishmania donovani*, a probable lipase. bioRxiv. 2022:2020.07. 23.218107.
- Zhang J, He J, Li J, Zhou Q, Chen H, Zheng Z et al. The immunogenicity and protective immunity of multi-epitopes DNA prime-protein boost vaccines encoding Amastin-Kmp-11, Kmp11-Gp63 and Amastin-Gp63 against visceral leishmaniasis. PloS one. 2020;15(3):e0230381.
- Stober CB, Lange UG, Roberts MT, Gilmartin B, Francis R, Almeida R et al. From genome to vaccines for leishmaniasis: screening 100 novel vaccine candidates against murine *Leishmania* major infection. Vaccine. 2006;24(14):2602-16.
- Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Molecular Biology and Evolution (submitted). 2015.
- McNicoll F, Müller M, Cloutier S, Boilard N, Rochette A, Dubé M et al. Distinct 3'-untranslated region elements regulate stage-specific mRNA accumulation and translation in *Leishmania*. Journal of Biological Chemistry. 2005;280(42):35238-46.
- Ribeiro PA, Vale DL, Dias DS, Lage DP, Mendonça DV, Ramos FF et al. *Leishmania infantum* amastin protein incorporated in distinct adjuvant systems induces protection against visceral leishmaniasis. Cytokine. 2020;129:155031.
- Lage DP, Vale DL, Linhares FP, Freitas CS, Machado AS, Cardoso JMO et al. A Recombinant Chimeric Protein-Based Vaccine Containing T-Cell Epitopes from Amastigote Proteins and Combined with Distinct Adjuvants, Induces Immunogenicity and Protection against *Leishmania infantum* Infection. Vaccines. 2022;10(7):1146.